

Multi-dimensional constraint relativistic mean field models and potential energy surfaces of actinide nuclei

Bing-Nan Lu,¹ Jie Zhao,¹ En-Guang Zhao,^{1,2} and Shan-Gui Zhou^{1,2,*}

¹*State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing 100190, China*

²*Center of Theoretical Nuclear Physics, National Laboratory of Heavy Ion Accelerator, Lanzhou 730000, China*
(Dated: April 10, 2013)

By breaking the reflection and the axial symmetries simultaneously, we developed multi-dimensional constraint relativistic mean field (MDC-RMF) models. In these models, the nuclear shape is assumed to be invariant under the reversion of x and y axes, i.e., the intrinsic symmetry group is V_4 and all shape degrees of freedom $\beta_{\lambda\mu}$ with even μ (β_{20} , β_{22} , β_{30} , β_{32} , β_{40} , ...) are included self-consistently. The RMF functional can be one of the following four forms: the meson exchange or point-coupling nucleon interactions combined with the non-linear or density-dependent couplings. The pairing effects are taken into account with the BCS approach. In this paper the formalism for MDC-RMF models is presented in details. Potential energy surface of ^{240}Pu is illustrated for numerical checks and for the study of the effects of triaxiality on the fission barriers. Potential energy curves of actinide nuclei around the first and second fission barriers are studied systematically with emphasis on the second ones which are lowered considerably by the triaxial deformation. We conclude that it is important to include the reflection asymmetric and non-axial shapes simultaneously for the study of potential energy surfaces and fission barriers of actinide nuclei and of those in unknown mass regions such as superheavy nuclei.

PACS numbers: 21.60.Jz, 24.75.+i, 25.85.-w, 27.90.+b

I. INTRODUCTION

The occurrence of spontaneous symmetry breaking leads to nuclear shapes with a variety of symmetries [1, 2]. A lot of nuclear phenomena are connected with the nuclear deformation, including the small and large amplitude collective motions, e.g., the rotation and the fission [3, 4]. There are also many unresolved problems concerning the motion or change along the nuclear shape degrees of freedom. For example, in recent years lots of efforts were devoted to explore nuclear shape phase transitions [5–8]; one way to investigate shape phase transitions is to examine nuclear potential energy surfaces and their changes with the nucleon number [9–13].

The shape of a nucleus can be described by the parametrization of the nuclear surface or the nucleon density distribution. There are mainly two kinds of ways to parametrize the nuclear shape [14, 15]. One of them is the two-center or two-center-like parametrization [16–20]. The other way is of one-center and to make a multipole expansion,

$$R(\theta, \varphi) = R_0 \left[1 + \beta_{00} + \sum_{\lambda=1}^{\infty} \sum_{\mu=-\lambda}^{\lambda} \beta_{\lambda\mu}^* Y_{\lambda\mu}(\theta, \varphi) \right]. \quad (1)$$

where $\beta_{\lambda\mu}$'s are deformation parameters. The way of multipole expansion is usually used in mean field calculations. The majority of observed nuclear shapes is

of spheroidal form which can be described by the axial-quadrupole deformation parameter β_{20} though in early years the Nilsson perturbed-spheroid parameter ϵ_2 was usually adopted for numerical convenience [21]. The non-axial-quadrupole (triaxial) deformation β_{22} (or, equivalently, γ) has been predicted in atomic nuclei for quite a long time and it may play important roles in superheavy nuclei [22]. A static triaxial shape manifests itself by the wobbling motion and chiral doublet bands; both have been extensively studied from experimental and theoretical sides [23–27]. Particularly, the existence of multiple chiral doublet (M χ D) bands in one nucleus was predicted [28]. Recently two distinct chiral doublet bands were observed in ^{133}Ce [29] which for the first time provides a convincing verification of the prediction. The softness with respect to the gamma distortion also results in many interesting nuclear phenomena [30–35].

The octupole shapes with $\lambda = 3$ are predicted to exist in nuclei in several mass regions [36]. The low-lying negative parity bands observed in actinides and some rare-earth nuclei are related to the reflection asymmetric shapes [37–43]. It was also found that the octupole deformation is responsible for the Ra puzzle [44], i.e., the anomalous enhancement of the residual proton-neutron interactions for Ra isotopes around $N = 135$ [45, 46]. The influences of the non-axial octupole β_{32} deformation on the low-lying spectra have been also traced out theoretically and experimentally [47–56]. For example, reflection asymmetric shell model calculations revealed that the observed low-energy 2^- bands in $N = 150$ nuclei [57] are caused by the β_{32} deformation [58]. Indeed, strong Y_{32} -correlations were found in some $N = 150$ isotones from multi-dimensional constraint covariant den-

* sgzhou@itp.ac.cn

sity functional theories (MDC-CDFT) [59]. Experimental efforts have also been made to explore the tetrahedral symmetry in nuclei [60, 61].

Deformations of higher multipoles with $\lambda > 3$ are important to different extents. The hexadecapole deformation, β_{40} or ϵ_4 , has been included in deformed mean field potentials since 1960s, see, e.g., Ref. [62]. The important effects of the high order deformation β_{60} or ϵ_6 on the angular momentum alignments and dynamic moments of inertia in superheavy nuclei are also discussed [63, 64].

The shape degrees of freedom are important not only for the ground states or small amplitude collective motions, but also for large amplitude collective motions such as fission. Since the discovery of the nuclear fission [65], the description of the fission process has been a difficult and challenging task. The fission dynamics are mostly determined by the barriers which prohibit the dissolving of the nucleus [66]. In order to study the fission problem, one should have very accurate information about the fission barrier, i.e., the height and width, or more precisely, the shape of the fission barrier [67, 68]. Particularly, to explore the island of stability of superheavy nuclei (SHN) [69–72], it is more and more desirable to have accurate predictions of fission barriers of SHN [73–81]. Until now many popular nuclear structure models have been employed to calculate the nuclear fission barriers, including the macroscopic-microscopic (MM) model [18, 73, 82–86], the extended Thomas-Fermi plus Strutinsky integral (ETFSI) method [87], the Hartree-Fock or Hartree-Fock-Bogoliubov method with the Skyrme force [88–93] and the Gogny force [94, 95], and the covariant density function theory [88, 96–102].

Besides β_{20} which describes the elongation of a fissile nucleus and β_{40} which is relevant to the size of a neck, many other shape degrees of freedom are also crucial for determining the shape of fission barriers and the fission path. Let's take actinides as examples. Due to the shell effects, actinide nuclei are characterized by a two-humped fission barrier [14] and there may also exist a third minimum [86, 103–107]. It has long been known from the MM model calculations that the inner fission barrier is lowered by the non-axial-quadrupole deformation [108–110] and the outer barrier by the reflection asymmetric (RA) shape [111]. Later, the important roles played by the non-axial-quadrupole deformation and octupole deformation were confirmed in the non-relativistic [112] and relativistic [113] density functional calculations, respectively. Therefore what is usually done is to consider the triaxial but reflection symmetric (RS) shapes for the inner barrier and axially symmetric but RA shapes for the outer one [73, 94, 114]. The non-axial octupole deformations are considered in both the MM models [115] and the non-relativistic Hartree-Fock theories [116].

In recent years, the nuclear CDFT has been very successful in the description of both the ground states and the exciting states of the nuclei ranging from light to superheavy regions [117–124]. Particularly, the CDFTs have been used to study the potential energy surfaces

(PES) and the fission barriers of heavy and superheavy nuclei [100, 101, 120, 125–128]. We have developed MDC-CDFTs by breaking the reflection and the axial symmetries simultaneously. In these theories, all shape degrees of freedom $\beta_{\lambda\mu}$ with even μ , e.g., β_{20} , β_{22} , β_{30} , β_{32} , β_{40} , $\dots$, are included self-consistently. The functional can be one of the following four forms: the meson exchange or point-coupling nucleon interactions combined with the non-linear or density-dependent couplings. For the particle-particle channel, either the BCS approach or the Bogoliubov transformation has been implemented and the pairing force could be δ -force or the finite-range separable force [129–131]. For convenience, the MDC-CDFT with the BCS approach for the pairing is named as MDC-RMF model and that with the Bogoliubov transformation as MDC-RHB one. Due to the heavy computational burden, MDC-RHB theories have not been used to do multi-dimensional constraint calculations yet (here “multi” means three or more). In Ref. [125, 126] we calculated the potential energy surfaces of actinide nuclei in the $(\beta_{20}, \beta_{22}, \beta_{30})$ deformation space with the MDC-RMF models and we found that the triaxiality also plays an important role upon the second fission barriers. In this paper, we will present the detailed formulae for MDC-RMF models and some results of actinide nuclei.

The paper is organized as follows. The formalism of the MDC-RMF models will be given in Sec. II. In Sec. III we present the numerical details and results on the PES's of the actinide nuclei. A summary is given in Sec. IV.

II. FORMALISM OF MULTI-DIMENSIONAL CONSTRAINED RELATIVISTIC MEAN FIELD MODELS

In the CDFTs a nucleus is treated as a composite of nucleons interacting through exchanges of mesons and photons. The effects of mesons are described either by the mean fields or by the point-like interactions between the nucleons. Meanwhile, the density dependence of the coupling constants [132, 133] or the non-linear coupling terms [134–136] are introduced to give correct saturation properties of the nuclear matter. Accordingly, the form of the covariant density functional can be one of the following four: the meson exchange or point-coupling nucleon interactions combined with the nonlinear or density dependent couplings. Most of the computational efforts are devoted to the solving of the Dirac equation and the calculation of the various densities which are common for all these CDFTs. In this section, we mainly present the formalism of the CDFT with the non-linear point couplings (NL-PC). For more details of CDFTs, the readers are referred to Refs. [117–124].

The starting point of the covariant density functional with NL-PC is the following Lagrangian:

$$\mathcal{L} = \bar{\psi}(i\gamma_\mu\partial^\mu - M)\psi - \mathcal{L}_{\text{lin}} - \mathcal{L}_{\text{nl}} - \mathcal{L}_{\text{der}} - \mathcal{L}_{\text{cou}}, \quad (2)$$

where

$$\begin{aligned}\mathcal{L}_{\text{lin}} &= \frac{1}{2}\alpha_S\rho_S^2 + \frac{1}{2}\alpha_V j_V^2 + \frac{1}{2}\alpha_{TS}\vec{\rho}_{TS}^2 + \frac{1}{2}\alpha_{TV}\vec{j}_{TV}^2, \\ \mathcal{L}_{\text{nl}} &= \frac{1}{3}\beta_S\rho_S^3 + \frac{1}{4}\gamma_S\rho_S^4 + \frac{1}{4}\gamma_V[j_V^2]^2, \\ \mathcal{L}_{\text{der}} &= \frac{1}{2}\delta_S[\partial_\nu\rho_S]^2 + \frac{1}{2}\delta_V[\partial_\nu j_V^\mu]^2 + \frac{1}{2}\delta_{TS}[\partial_\nu\vec{\rho}_{TS}]^2 \\ &\quad + \frac{1}{2}\delta_{TV}[\partial_\nu\vec{j}_{TV}]^2, \\ \mathcal{L}_{\text{cou}} &= \frac{1}{4}F^{\mu\nu}F_{\mu\nu} + e\frac{1-\tau_3}{2}A_\mu j_V^\mu,\end{aligned}\quad (3)$$

are the linear coupling, non-linear coupling, derivative coupling, and the Coulomb part, respectively. M is the nucleon mass, α_S , α_V , α_{TS} , α_{TV} , β_S , γ_S , γ_V , δ_S , δ_V , δ_{TS} , and δ_{TV} are coupling constants for different channels and e is the electric charge. ρ_S , $\vec{\rho}_{TS}$, j_V , and $\vec{j}_{TV}$ are the iso-scalar density, iso-vector density, iso-scalar current, and iso-vector current, respectively. The various densities and currents are defined as:

$$\begin{aligned}\rho_S &= \bar{\psi}\psi, & \vec{\rho}_{TS} &= \bar{\psi}\vec{\tau}\psi, \\ j_V^\mu &= \bar{\psi}\gamma^\mu\psi, & \vec{j}_{TV}^\mu &= \bar{\psi}\vec{\tau}\gamma^\mu\psi.\end{aligned}\quad (4)$$

Starting from the Lagrangian, using the Slater determinants as trial wave functions, and neglecting the Fock term as well as the contributions to the densities and currents from the Dirac sea, we can derive the relativistic mean field equations with the variational principle:

$$\hat{h}\psi_k(\mathbf{r}) = \epsilon_k\psi_k(\mathbf{r}), \quad (5)$$

where

$$\hat{h} = \boldsymbol{\alpha} \cdot [\mathbf{p} - \mathbf{V}(\mathbf{r})] + \beta [M + S(\mathbf{r})] + V_0(\mathbf{r}), \quad (6)$$

is the single particle Hamiltonian and

$$S = \alpha_S\rho_S + \alpha_{TS}\vec{\rho}_{TS} \cdot \vec{\tau} + \beta_S\rho_S^2 + \gamma_S\rho_S^3 \\ + \delta_S\Delta\rho_S + \delta_{TS}\Delta\vec{\rho}_{TS} \cdot \vec{\tau}, \quad (7)$$

$$V^\mu = \alpha_V j_V^\mu + \alpha_{TV}\vec{j}_{TV}^\mu \cdot \vec{\tau} + \gamma_V j_V^2 j_V^\mu \\ + \delta_V\Delta j_V^\mu + \delta_{TV}\Delta\vec{j}_{TV}^\mu \cdot \vec{\tau}, \quad (8)$$

are the scalar and vector potentials, respectively. In the present work we suppose that the states are invariant under the time-reversal operation, which means that all the time-odd or the vector components of the currents and the potentials vanish. In this case the single particle Hamiltonian has the time-reversal symmetry which simplifies the calculations. For convenience two vector densities are defined as the time-like components of the 4-currents j_V and $\vec{j}_{TV}$:

$$\rho_V(\mathbf{r}) = j_V^0(\mathbf{r}), \quad \rho_{TV}(\mathbf{r}) = j_{TV}^0(\mathbf{r}) \quad (9)$$

which are the only non-vanishing components.

It is customary to solve the deformed CDFT by expanding the axillary single particle wave functions on

a group of complete basis, e.g., the harmonic oscillator (HO) basis [137, 138] or the Woods-Saxon (WS) basis [139–143]. A reflection asymmetric relativistic mean field (RAS-RMF) approach has developed by using a basis in a two-centre harmonic-oscillator potential [144]. In our MDC-CDFTs, the single particle wave functions and various densities are expanded on an axially deformed harmonic oscillator (ADHO) basis [137, 138, 145]. The ADHO basis are defined as the eigen solutions of the Schrödinger equation,

$$\left[-\frac{\hbar^2}{2M}\nabla^2 + V_B(z, \rho)\right]\Phi_\alpha(\mathbf{r}\sigma) = E_\alpha\Phi_\alpha(\mathbf{r}\sigma), \quad (10)$$

where

$$V_B(z, \rho) = \frac{1}{2}M(\omega_\rho^2\rho^2 + \omega_z^2z^2), \quad (11)$$

is the ADHO potential and ω_z and ω_ρ are the oscillator frequencies along and perpendicular to the symmetry z axis, respectively. The solution of Eq. (10) reads

$$\Phi_\alpha(\mathbf{r}\sigma) = C_\alpha\phi_{n_z}(z)R_{n_\rho}^{m_l}(\rho)\frac{1}{\sqrt{2\pi}}e^{im_l\varphi}\chi_{s_z}(\sigma), \quad (12)$$

where $\phi_{n_z}(z)$ and $R_{n_\rho}^{m_l}(\rho)$ are the HO wave functions,

$$\begin{aligned}\phi_{n_z}(z) &= \frac{1}{\sqrt{b_z}}\frac{1}{\pi^{1/4}\sqrt{2^{n_z}n_z!}}H_{n_z}\left(\frac{z}{b_z}\right)e^{-\frac{z^2}{2b_z^2}}, \\ R_{n_\rho}^{m_l}(\rho) &= \frac{1}{b_\rho}\sqrt{\frac{2n_\rho!}{(n_\rho + |m_l|)!}}\left(\frac{\rho}{b_\rho}\right)^{|m_l|}L_{n_\rho}^{|m_l|}\left(\frac{\rho^2}{b_\rho^2}\right)e^{-\frac{\rho^2}{2b_\rho^2}}.\end{aligned}\quad (13)$$

χ_{s_z} is a two component spinor and C_α is a complex number introduced for convenience. Harmonic oscillator lengths b_z and b_ρ are related to the frequencies by $b_z = 1/\sqrt{M\omega_z}$ and $b_\rho = 1/\sqrt{M\omega_\rho}$. The corresponding eigen energy is $E_\alpha = \omega_\rho(2n_\rho + |m_l| + 1) + \omega_z(n_z + \frac{1}{2})$ and the major quantum number is $N_\alpha = 2n_\rho + |m_l| + n_z$.

These basis are also eigen functions of the z component of the angular momentum, $\hat{j}_z$ with eigen values $K = m_l + m_s$. For any basis $\Phi_\alpha(\mathbf{r}\sigma)$ the time reversed basis is defined as $\Phi_{\bar{\alpha}}(\mathbf{r}\sigma) = \mathcal{T}\Phi_\alpha(\mathbf{r}\sigma)$, where $\mathcal{T} = i\sigma_y K$ is the time reversal operator and K is the complex conjugation. Apparently we have $K_{\bar{\alpha}} = -K_\alpha$ and $\pi_{\bar{\alpha}} = \pi_\alpha$ where $\pi = \pm 1$ is the parity. The deformation of the basis β_{basis} is defined through the relations $\omega_z = \omega_0 \exp\left(-\sqrt{\frac{5}{4\pi}}\beta_{\text{basis}}\right)$ and $\omega_\rho = \omega_0 \exp\left(\sqrt{\frac{5}{16\pi}}\beta_{\text{basis}}\right)$, where $\omega_0 = (\omega_z\omega_\rho^2)^{1/3}$ is the frequency of the corresponding spherical oscillator [137, 138].

These basis form a complete set for expanding any two component spinors. For a Dirac spinor with four components,

$$\psi_i(\mathbf{r}\sigma) = \left(\sum_\alpha f_i^\alpha\Phi_\alpha(\mathbf{r}\sigma)\right), \quad (14)$$

where the sum runs over all the possible combination of the quantum numbers $\alpha = \{n_z, n_r, m_l, m_s\}$, f_i^α and

g_i^α are the expansion coefficients. In practical calculations the sum in Eq. (14) are truncated as $n_z/Q_z + (2n_\rho + |m|)/Q_\rho \leq N_{\text{ph}}$, where N_{ph} is an integer and $Q_z = \max(1, b_z/b_0)$ and $Q_\rho = \max(1, b_\rho/b_0)$ are constants related to the oscillator lengths b_0 , b_z , and b_ρ . Clearly in any case the number of the basis included are no less than that obtained from the major shell truncation $N \leq N_{\text{ph}}$.

If a nucleus is invariant under the rotation around z axis and the spatial reflection, the angular momentum projection and the parity are conserved and the RMF equation (5) can be decomposed into blocks characterized by the quantum numbers K and π . Usually only half of the space with $K_\alpha > 0$ is considered due to the time-reversal symmetry.

Now let us turn to the general case that the axial symmetry as well as the spatial reflection symmetry are broken. If we still expand the spinors on the ADHO basis, different components with different K and π are mixed together, that is, we must diagonalize a larger single particle Hamiltonian matrix with non-zero matrix elements between two basis states with different K and π . Nevertheless, even in this case we still can have one symmetry operator that makes the Hamiltonian matrix block-diagonal. Due to the axial symmetry of the basis, it is convenient to choose the simplex operator $\hat{S} = ie^{-i\pi j_z}$ as a conserved quantity. Note that for a fermion system with a half-integer spin, $\hat{S}$ is a Hermitian operator and $\hat{S}^2 = 1$. This operator corresponds to the rotation by π around the z axis thus leaving the nuclear mean field invariant. S is also a good quantum number for the basis with $S\Phi_\alpha = (-1)^{K_\alpha-1/2}\Phi_\alpha$, which means that the basis Φ_α with $K_\alpha = +1/2, -3/2, 5/2, -7/2, \dots$ span the subspace with $S = 1$, while their time-reversed states span the one with $S = -1$. Note that now the blocks with $K = +1/2$ and $K = -3/2$ (not that of $K = +3/2$) are mixed. Remembering that for a nucleus with the time reversal symmetry only the basis with $S = 1$ are used in the expansions, for such basis we set $C_\alpha = 1$ when expanding the large components and $C_\alpha = i$ for the small ones. The basis with $S_\alpha = -1$ are obtained by simply applying $\mathcal{T}$ on them. Furthermore, for systems with the time-reversal symmetry, it is only necessary to diagonalize the matrix with $S = 1$ and the other half is obtained by a time reversal operation on the obtained single particle wave functions.

For deformed nuclei with the V_4 symmetry, we expand the potentials $V(\mathbf{r})$, $S(\mathbf{r})$, and the densities in Eq. (4) in terms of the Fourier series,

$$f(\rho, \varphi, z) = \sum_{\mu=-\infty}^{\infty} f_\mu(\rho, z) \frac{1}{\sqrt{2\pi}} \exp(i\mu\varphi). \quad (15)$$

By substituting this expansion into the symmetry conditions, it is easy to see that $f_\mu = f_\mu^* = f_{-\mu}$ and $f_n = 0$ for

odd n . Thus the expansion (15) can be simplified as

$$f(\rho, \varphi, z) = f_0(\rho, z) \frac{1}{\sqrt{2\pi}} + \sum_{n=1}^{\infty} f_n(\rho, z) \frac{1}{\sqrt{\pi}} \cos(2n\varphi), \quad (16)$$

where

$$\begin{aligned} f_0(\rho, z) &= \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} d\varphi f(\rho, \varphi, z), \\ f_n(\rho, z) &= \frac{1}{\sqrt{\pi}} \int_0^{2\pi} d\varphi f(\rho, \varphi, z) \cos(2n\varphi), \end{aligned} \quad (17)$$

are real functions of ρ and z . The details for calculating the matrix elements of the Dirac Hamiltonian and various densities and their derivatives are given in the Appendix.

For open shell nuclei the pairing interaction becomes crucial and must be included. For example, the calculated fission barriers depend very much on the form and strength of the effective pairing interactions [146]. In realistic calculations several methods have been used to treat the pairing effects, e.g., the BCS approach, the Bogoliubov transformation, and the particle number conserving method [147–151]. Here we show the formulas for the BCS approximation with a δ pairing force which are incorporated in the MDC-RMF models. In the particle-particle channel, the gap equation reads [2],

$$\Delta_k = \sum_{k'>0} \frac{1}{2} V_{kkk'\bar{k}'}^{\text{pp}} \frac{\Delta_{k'}}{\sqrt{\tilde{\epsilon}_{k'}^2 + \Delta_{k'}^2}}, \quad (18)$$

where $\tilde{\epsilon}_k = \epsilon_k - \lambda$ and λ is the Fermi energy. The total pairing energy is

$$E_{\text{pair}} = \frac{1}{4} V_0 \int d^3\mathbf{r} \kappa^*(\mathbf{r}) \kappa(\mathbf{r}). \quad (19)$$

To obtain the PES one can perform a constraint calculation which is equivalent to adding an external potential during the iteration [2]. The quadratic constraint method is usually used,

$$E' = E_{\text{RMF}} + \sum_{\lambda\mu} \frac{1}{2} C_{\lambda\mu} (Q_{\lambda\mu} - m_{\lambda\mu})^2, \quad (20)$$

where $C_{\lambda\mu}$ is the spring constant and $m_{\lambda\mu}$'s are desired moments. With this method the calculation always converges to a deformation point on the PES other than the desired one. To overcome this shortcoming and to get a PES with equally distributed points, we use a modified linear constraint method. The Routhian reads,

$$E' = E_{\text{RMF}} + \sum_{\lambda\mu} \frac{1}{2} C_{\lambda\mu} Q_{\lambda\mu}, \quad (21)$$

with the variables $C_{\lambda\mu}$'s changing their values during the iteration through the following relation,

$$C_{\lambda\mu}^{(n+1)} = C_{\lambda\mu}^{(n)} + k_{\lambda\mu} (\beta_{\lambda\mu}^{(n)} - \beta_{\lambda\mu}), \quad (22)$$

where $\beta_{\lambda\mu}$ is the desired deformation, $k_{\lambda\mu}$ is a constant, and $C_{\lambda\mu}^{(n)}$ is the value at the n th step. We found that this constraint method works well in our multi-dimensional calculations.

The total energy of the nucleus reads

$$E_{\text{total}} = \int d^3\mathbf{r} \left\{ \sum_k v_k^2 \psi_k^\dagger (\boldsymbol{\alpha} \cdot \mathbf{p} + \beta M) \psi_k + \frac{1}{2} \alpha_S \rho_S^2 + \frac{1}{2} \alpha_V \rho_V^2 + \frac{1}{2} \alpha_{TS} \rho_{TS}^2 + \frac{1}{2} \alpha_{TV} \rho_{TV}^2 + \frac{1}{3} \beta_S \rho_S^3 + \frac{1}{4} \gamma_S \rho_S^4 + \frac{1}{4} \gamma_V \rho_V^4 + \frac{1}{2} \delta_S \rho_S \Delta \rho_S + \frac{1}{2} \delta_V \rho_V \Delta \rho_V + \frac{1}{2} \delta_{TS} \rho_{TS} \Delta \rho_{TS} + \frac{1}{2} \delta_{TV} \rho_{TV} \Delta \rho_{TV} + \frac{1}{2} e \rho_C A \right\} + E_{\text{pair}} + E_{\text{c.m.}}, \quad (23)$$

where $E_{\text{c.m.}}$ is the center of mass correction which can be calculated either from the quasi-particle vacuum,

$$E_{\text{c.m.}} = -\frac{\langle \hat{P}^2 \rangle}{2MA}, \quad (24)$$

where $\hat{P}$ is the total linear momentum and A is the nuclear mass number or in the oscillator approximation,

$$E_{\text{c.m.}} = -\frac{3}{4} \times 41 \times A^{1/3} \text{ MeV}, \quad (25)$$

depending on the effective interactions used in the RMF functional.

The intrinsic multipole moments are calculated from the densities by

$$Q_{\lambda\mu} = \int d^3\mathbf{r} \rho_V(\mathbf{r}) r^\lambda Y_{\lambda\mu}(\Omega), \quad (26)$$

where $Y_{\lambda\mu}(\Omega)$ is the spherical harmonics. The deformation parameter $\beta_{\lambda\mu}$ is obtained from the corresponding multipole moment by

$$\beta_{\lambda\mu} = \frac{4\pi}{3NR^\lambda} Q_{\lambda\mu}, \quad (27)$$

where R is the estimated nuclear radius $R = 1.2 \times A^{1/3}$ fm and N is the number of protons, neutrons, or nucleons.

III. RESULTS AND DISCUSSIONS

A. Numerical details

In this section we present the numerical details and some illustrative calculations of MDC-RMF models. The potentials and densities are calculated on a spatial lattice of which mesh points in the ρ and z directions are designed in a way that the Gaussian quadrature can be




FIG. 1. (Color online) The potential energy curve of ^{240}Pu calculated with different number of the oscillator shells N_{ph} from the MDC-CDFT calculations. The axial symmetry is imposed. The results calculated with $N_{\text{ph}} = 16, 18$, and 20 are depicted by solid, dashed, and dotted lines, respectively. The four sub-figures show the detailed structure of the PES's near the ground state, the inner barrier, the isomeric state, and the outer barrier. The width of each sub-figure is 0.1 and the height is 1 MeV.

made and those for the azimuthal angle are equally distributed. Since we keep the mirror reflection symmetry with respect to the $x = 0$ or $y = 0$ planes, only mesh points with positive x and y are considered. For the azimuthal angle, more than 10 mesh points for light nuclei and about 20 mesh points for heavy nuclei are used, which are enough for most of practical applications. In two special cases the number of the mesh points can be even reduced: (1) for axial symmetric nuclei the azimuthal degree of freedom vanishes and (2) for reflection symmetric nuclei the mesh points with $z < 0$ can be omitted. With these choices the values of the localized fields and potentials in the full lattice space can be simply obtained by symmetry transformations like the rotations or space reflection. Nevertheless, the Coulomb field must be treated carefully due to its long range nature.

The calculated physical observables should converge as the truncation $N_{\text{ph}} \rightarrow \infty$. In Fig. 1 we show the potential energy curve of ^{240}Pu calculated with different truncations, $N_{\text{ph}} = 16, 18$, and 20 . The effective interaction PC-PK1 [152] is used. For simplicity the calculations are restricted to the axially symmetric case. As is found in most of earlier calculations, the results show a typi-




FIG. 2. (Color online) Calculated total binding energy of ^{240}Pu by using basis with different basis deformation β_{basis} and different number of major shells N_{ph} . The results calculated with 16, 18, and 20 shells are denoted by red, green, and blue dots, respectively. The deformation is constrained to $\beta_{20} = 0.3, 1.3, 2.0$, and 3.0 , respectively.

cal two-humped structure and the reflection asymmetric shapes are favored in the outer barrier region. To see more clearly the truncation errors we have amplified the figure near four important points, i.e., the ground state, the top of the inner barrier, the isomeric state, and the top of the outer barrier. From the four sub-figures we can investigate the convergence properties of our model. When N_{ph} increases from 16 to 20, the binding energy changes less than 200 keV for the ground state and less than 150 keV for the second minimum. The changes of the first and the second barrier heights are within 200 keV and 300 keV, respectively.

Next we show how the results depend on the deformation of the ADHO potential which is used to generate the ADHO basis; for brevity, we will call it the basis deformation and label it with β_{basis} . In Fig. 2 we depict the calculated binding energy as a function of the basis deformation. In principle, if the basis space is complete or nearly complete, the results should not change a lot when the basis deformation changes. Near the ground state $\beta_{20} = 0.3$, the calculated energies are rather stable

against the basis deformation. Furthermore, the results with $N_{\text{ph}} = 16, 18$, and 20 coincide with each other. This conclusion holds also for the second barrier, $\beta_{20} = 1.3$, except that two points with small basis deformations and $N_{\text{ph}} = 16$ are very high. However, if we choose properly the basis deformation, 16 shells are enough for calculations around the barriers. For even larger deformation, $\beta_{20} = 2.0$ or 3.0 , the results for $N_{\text{ph}} = 16$ deviate from that for $N_{\text{ph}} = 18$ and 20 ; the differences between the results for $N_{\text{ph}} = 18$ and 20 are still small.

When the basis is not complete, the calculated single particle energies contain the contributions from the other levels in both the Fermi sea and the Dirac sea. The contributions from the Fermi sea raises the total energy, while that from the Dirac sea lowers the energy. Thus as more basis states are included in the expansion, the binding energy may not increase monotonically. This is different from the non-relativistic calculations where the calculated binding energy always increases when the basis size is increased. In the whole deformation range considered here, the largest difference between the results with $N_{\text{ph}} = 16$ and $N_{\text{ph}} = 20$ appears at the top of the second barrier. This difference is about 300 keV. It is very time consuming if non-axial deformations are included in the calculations. In the present work, for the PES's of actinide nuclei we use $N_{\text{ph}} = 20$ in axially symmetric calculations and $N_{\text{ph}} = 16$ when the triaxial deformations are included.

B. Three-dimensional PES of ^{240}Pu

The double-humped fission barriers of actinide nuclei are usually used as a benchmark for theoretical models [91, 96, 98, 99, 112, 113, 153–160]. As a nucleus evolves from the ground state to the fission configurations, various shape degrees of freedom play important and different roles in determining the heights of the inner and outer barriers. It has long been known that the inner barrier is lowered when the triaxial deformation is allowed, while for the outer barrier, the reflection asymmetric shape is favored. We have shown that the reflection asymmetric outer barrier may be further lowered by including the non-axial shape degrees of freedom [125–127]. In this section we show some results of multi-dimensional PES's for actinides. We will first present detailed results about the three dimensional PES of ^{240}Pu , then display systematic results of the actinide nuclei. In these calculations, the effective interaction PC-PK1 [152] is used.

In Figs. 3 and 4 is shown the calculated three dimensional PES of ^{240}Pu . In each sub-figure we fix the value of β_{20} and display the energy as a function of β_{22} and β_{30} . In other words, we have made three-dimensional constraints on the corresponding multipole moments. Note that other shape degrees of freedom $\beta_{\lambda\mu}$ with even μ , e.g., $\beta_{32}, \beta_{40}, \beta_{42}, \beta_{44}, \beta_{50}, \dots$, are also included in the calculations self-consistently. The MDC-RMF equations are solved for each point on the deformation lattice




FIG. 3. (Color online) Sections of the three-dimensional potential energy surfaces $E = E(\beta_{20}, \beta_{22}, \beta_{30})$ of ^{240}Pu near the inner barrier from MDC-RMF calculations. In each sub-figure the energy is shown as a function of the deformation parameters β_{22} and β_{30} when β_{20} is fixed to certain values. The energies are normalized with respect to the binding energy of the ground state. The contour interval is 1 MeV.




FIG. 4. (Color online) Sections of the three-dimensional potential energy surfaces $E = E(\beta_{20}, \beta_{22}, \beta_{30})$ of ^{240}Pu near the outer barrier from MDC-RMF calculations. In each sub-figure the energy is shown as a function of the deformation parameters β_{22} and β_{30} when β_{20} is fixed to certain values. The energies are normalized with respect to the binding energy of the ground state. The contour interval is 1 MeV.

$(\beta_{20}, \beta_{22}, \beta_{30})$ in which β_{20} runs from 0.25 to 0.95 with a step size of 0.05, β_{22} from 0 to 0.25 with a step size of 0.01, and β_{30} from 0 to 0.50 with a step size of 0.05. The points with $\beta_{22}(\beta_{30}) < 0$ are obtained through the relation $E(\beta_{20}, \beta_{22}, \beta_{30}) = E(\beta_{20}, |\beta_{22}|, |\beta_{30}|)$. That is, for plotting each sub-figure 26 (for β_{22}) $\times$ 11 (for β_{30}) = 286 points are calculated, and totally 286×15 (for β_{20}) = 4290 points are calculated for Figs. 3 and 4. This deformation lattice covers the shape space of the most interest for ^{240}Pu , from the ground state to the isomeric state and fissioning configurations.

Now let us examine the first three sub-figures with $\beta_{20} = 0.25, 0.30$, and 0.35 . It is clear that the ground state with $\beta_{20} \sim 0.3$ is both axial symmetric and reflection symmetric, though it is a little soft against the octupole distortion. When the nucleus is stretched by the quadrupole constraining potential, it becomes softer against the triaxial as well as the octupole distortions. From the sub-figures with $\beta_{20} = 0.40 \sim 0.65$ one finds that two symmetric minima with non-zero triaxial deformation β_{22} appear and the corresponding fission paths are much more favored than the axial symmetric one, which is consistent with earlier calculations [100]. Although $\beta_{30} = 0$ for all minima in these sub-figures, the softness against the octupole distortion changes as the nucleus is elongated. The inner fission barrier locates near the deformation $\beta_{20} \sim 0.60$. For the last five sub-figures with $\beta_{20} = 0.70 \sim 0.95$, the situation becomes more simpler, where the nucleus becomes axial symmetric and reflection symmetric again. Nevertheless, the trend to become softer against the octupole distortion can be seen from these sub-figures, which indicates that the reflection asymmetric shape becomes more relevant.

As the deformation β_{20} becomes larger, the second minimum and the second saddle point of the PES appear. From the last three sub-figures of Fig. 3 and the first three sub-figures of Fig. 4 we see that, the nucleus keeps reflection symmetric near the fission isomeric state with $\beta_{20} \sim 0.95$, but it becomes softer against the β_{30} distortion. Note that the scales of the β_{30} axes are different in Figs. 3 and 4. At $\beta_{20} = 1.15$, two minima corresponding to reflection asymmetric shapes appear. Here the effects of the non-axial deformation is not apparent, but along the β_{22} direction the PES becomes softer. Finally, at $\beta_{20} = 1.2$ the energy of the reflection asymmetric shape is lower by about 1 MeV than that of the reflection symmetric one. Interestingly, when β_{20} increases continuously, beyond the top of the second barrier with $\beta_{20} = 1.2$, each reflection asymmetric minimum splits into two minima with non-zero β_{22} . The energy gain due to the triaxial distortion is about 1 MeV. The nucleus becomes axial symmetric again when $\beta_{20} > 1.4$.

From above discussions, we can draw the following conclusions for ^{240}Pu : (1) Both the ground state and the fission isomeric state are axial and reflection symmetric; (2) Around the first fission barrier it assumes triaxial and reflection symmetric shapes; (3) Around the second fission barrier both triaxial and octupole deformations are

important.

C. PES's of even-even actinide nuclei around fission barriers

The self-consistent three-dimensional constraint calculations are very time-consuming, we have only performed such calculations for ^{240}Pu . From this benchmark study we learned many experiences about the important roles played by various shape degrees of freedom in different regions of the deformation space, including the conclusions listed in the end of Sec. III B. These experiences are used in a systematic study of even-even actinide nuclei.

Since around the inner barrier an actinide nucleus assumes triaxial but reflection symmetric shapes, we can make a one-dimensional constraint calculation with the triaxial deformation allowed and the reflection symmetry imposed. In Fig. 5 we present potential energy curves of even-even actinide nuclei around the first fission barrier from MDC-RMF calculations. The axial symmetric results are displayed by dashed lines and those from triaxial calculations are shown by solid lines. The empirical values of fission barrier heights are taken from Ref. [161] and labelled with solid red circles. We also list in Table I the heights of the first and second fission barriers of actinide nuclei from MDC-RMF calculations compared with the empirical values. It is clearly seen that the triaxial deformation lowers the inner barrier of these actinide nuclei by about 1 $\sim$ 4 MeV and this is consistent with Ref. [100]. The agreement of our calculation results with the empirical ones is good for most of the nuclei studied here with exceptions in the two thorium isotopes and $^{238,240}\text{U}$. We have discussed possible reasons for these discrepancies in Ref. [125].

It is more complicated to calculate the second fission barrier height because more shape degrees of freedom become important and there are often two or more fission paths. What we have done is the following [125]: (1) The axial symmetry is assumed and a two-dimensional constraint calculation in the (β_{20}, β_{30}) plane is made; (2) From the two-dimensional calculation the lowest fission path $\beta_{30}^{\text{lowest}}(\beta_{20})$ is approximately identified; (3) Along this fission path, a one-dimensional β_{20} -constraint calculation is performed with the triaxial and octupole deformations allowed; (4) At each point with β_{20} fixed, the initial deformations are taken as $\beta_{22}^{\text{ini}} = 0$ and $\beta_{30}^{\text{ini}} = \beta_{30}^{\text{lowest}}(\beta_{20})$; (5) In this one-dimensional potential energy curve, the second saddle point is located and the second barrier height is extracted.

Potential energy curves of even-even actinide nuclei around the second fission barrier from MDC-RMF calculations are shown in Fig. 6 from which and Table I one finds that for most of the nuclei investigated here, the triaxiality lowers the second barrier by about 0.5 $\sim$ 1 MeV, accounting for about 10 $\sim$ 20% of the barrier height. The calculation results with the triaxiality included agrees well with the empirical values for all ac-




FIG. 5. (Color online) Potential energy curves of even-even actinide nuclei around the first fission barrier from MDC-RMF calculations. The axial symmetric results are displayed by dashed lines, while those from the triaxial calculations are shown by solid lines. The binding energies are normalized with respect to the binding energies of the ground states of each nucleus. The empirical values of fission barriers are taken from Ref. [161].




FIG. 6. (Color online) Potential energy curves of even-even actinide nuclei around the second fission barrier from MDC-RMF calculations. The reflection asymmetric shapes are allowed in the calculations. The axial symmetric results are displayed by dashed lines, while those from the triaxial calculations are shown by solid lines. The binding energies are normalized with respect to the binding energies of the ground states of each nucleus. The empirical values of fission barriers are taken from Ref. [161].

TABLE I. Heights of the first and second fission barriers of some even-even actinide nuclei from MDC-RMF calculations compared with the empirical values (“Emp”) which are taken from Ref. [161]. The calculation results with and without the axial symmetry imposed are denoted by “AS” (axial symmetric) and “TA” (triaxial), respectively. The height is in MeV.

Nucleus	Z	N	A	First barrier			Second barrier		
				AS	TA	Emp	AS	TA	Emp
²³⁰ Th	90	140	230	5.03	3.96	6.10	6.80	6.37	6.50
²³² Th	90	142	232	4.94	4.12	5.80	6.70	6.18	6.70
²³² U	92	140	232	5.71	4.81	4.90	6.20	5.64	5.40
²³⁴ U	92	142	234	6.15	5.09	4.80	6.20	5.55	5.50
²³⁶ U	92	144	236	6.40	5.11	5.00	6.15	5.31	5.70
²³⁸ U	92	146	238	6.54	5.03	6.30	6.20	5.42	5.50
²⁴⁰ U	92	148	240	6.58	4.96	6.10	6.38	5.43	5.80
²³⁸ Pu	94	144	238	7.72	5.96	5.60	6.05	5.56	5.10
²⁴⁰ Pu	94	146	240	7.98	5.92	6.05	6.24	5.60	5.20
²⁴² Pu	94	148	242	8.05	5.77	5.85	6.43	5.74	5.10
²⁴⁴ Pu	94	150	244	7.85	5.40	5.70	6.26	5.49	4.90
²⁴⁶ Pu	94	152	246	7.37	4.76	5.40	5.84	4.96	5.30
²⁴² Cm	96	146	242	8.80	6.49	6.65	5.72	4.85	5.00
²⁴⁴ Cm	96	148	244	9.04	6.34	6.18	5.90	4.88	5.10
²⁴⁶ Cm	96	150	246	8.89	5.84	6.00	5.40	4.62	4.80
²⁴⁸ Cm	96	152	248	8.43	5.35	5.80	4.10	—	4.80
²⁵⁰ Cm	96	154	250	7.77	4.79	5.40	2.60	—	4.40
²⁵⁰ Cf	98	152	250	8.87	5.70	5.60	2.40	—	3.80
²⁵² Cf	98	154	252	8.41	5.26	5.30	1.20	—	3.50

tinide nuclei shown in Fig. 6. In Ref. [125], we have found that for ²⁴⁸Cm, the height of the second barrier without the triaxial deformation included is already smaller than the empirical value. From Table I it is seen that this is also the case for ²⁵⁰Cm and ^{250,252}Cf. Therefore the heights of the second barriers with the triaxial deformation included are not listed for these nuclei. The reason for these discrepancies may be related to a strong competition between the two or among more fission paths and the assumption on the barrier width made when the empirical value of the barrier height is evaluated [125].

From Figs. 5 and 6 one can find some interesting features concerning the triaxial shape of actinides around the first and the second barriers. Though it is emphasized here and in Refs. [100, 101] that the triaxiality is crucial in determining the height of these barriers, it is not necessary that the nucleus in question is triaxially deformed at these two saddle points. One can find that all of these even-even actinide nuclei shown in Fig. 5 are axially deformed at the first saddle points from our MDC-RMF calculations. From Fig. 6 it is seen that except ^{232,234,240}U, all other nuclei are also axially deformed at the second saddle points. The triaxial shape is only important before the saddle point around the first barriers, while it is often so after the saddle point around the second barrier, with exceptions in uranium isotopes. Therefore the two saddle points become closer when the triaxial deformation is included. Further investigations should be made to reveal the physics behind these features.

TABLE II. Heights of the second barrier of ²³⁴U calculated from MDC-RMF models with different parameter sets. The results with and without non-axial deformations included and their difference are denoted by B_{AS} , B_{TA} , and ΔB , respectively.

Parameter set	B_{AS} (MeV)	B_{TA} (MeV)	ΔB (MeV)
NL3*	7.54	6.85	0.69
NL-Z2	4.83	3.91	0.92
PC-PK1	6.20	5.55	0.65
DD-ME2	8.19	7.51	0.68
DD-PC1	6.13	5.64	0.49

D. Parameter dependence of the effects of triaxiality on the second barrier height

From the above detailed studies of the PES’s of actinide nuclei in the $(\beta_{20}, \beta_{22}, \beta_{30})$ space, it is clear that the triaxiality plays an important role upon the second fission barriers of actinide nuclei. We have studied the parameter dependence of the effects of triaxiality on the second barrier height and some results can be found in Refs. [125, 126, 162]. In Table II are listed the second barrier heights of ²³⁴U calculated with different parameter sets, including meson-exchange ones NL3* [163], NL-Z2 [164], DD-ME2 [165], and point-coupling ones PC-PK1 [152] and DD-PC1 [166]. One finds that the second barrier height may differ by a few MeV with different effective interactions, but in all cases the barriers are lowered considerably by the non-axial deformations. For this specific nucleus, the lowering effects with different effective interactions are about at least 0.5 MeV and may as large as 0.9 MeV which are clearly larger than the possible errors from the basis truncations discussed in Sec. III A.

E. Further discussions

A systematic study of fission barriers is performed for even-even superheavy nuclei with $Z = 112 \sim 120$ by using covariant density functional models and the outer fission barriers are found to be considerably affected by the triaxiality [101]. However, in a recent work within the macroscopic-microscopic model, it was found that the influence of the triaxiality on second fission barriers is negligible [167]. This raises an open problem: What are the main reasons for the different effects of the triaxiality on the second fission barriers predicted by the MM model and CDFTs? It would be interesting to make more investigations with different models, especially, with the non-relativistic DFTs.

One of the problems concerning the PES calculated from self-consistent approaches is that there may exist unexpected discontinuities on the PES’s [73]. This is mainly due to the complexity of multi-dimensional PES’s. When a high-dimensional PES is projected onto a low-dimensional one, the minimizing procedure incorporated

implicitly in the self-consistent approaches may cause a sudden change in the total energy at some points in the low-dimensional deformation space. This may result in spurious saddle points. In our MDC-RMF calculations, we have tried some ways to avoid discontinuities on the PES's and to exclude spurious saddle points [125]. To be completely free of discontinuities or spurious saddle points, one certainly should carry out multi-dimensional constraint calculations with higher-polarity deformations included. In Ref. [168] the authors analyzed the origin of the discontinuities and proposed a numerical method to identify them. It will be useful to implement this method in the present MDC-RMF calculations; this will be one of our topics in the future.

Since many symmetries are broken in mean field calculations, quantum numbers related to these symmetries are not conserved any more [2]. For example, the total angular momentum or nuclear spin J is not a good quantum number when β_{20} is not zero; similarly, the projection of the total angular momentum on the symmetry axis K is also not conserved if a nucleus is triaxially deformed [169]. When the reflection symmetry is broken, the parity is not a good quantum number. In our three-dimensional constraint calculations, all these quantum numbers are not good ones. One may introduce techniques of projection to restore the broken symmetries [2]. In recent years, the importance of various projections on the PES's and fission barriers have been investigated [94, 160, 170, 171].

IV. SUMMARY

We developed multi-dimensional constraint relativistic mean field (MDC-RMF) models. In these models, the nuclear shape is assumed to be invariant under the reversion of x and y axes, i.e., the intrinsic symmetry group is V_4 and all shape degrees of freedom $\beta_{\lambda\mu}$ with even μ (β_{20} , β_{22} , β_{30} , β_{32} , β_{40} , $\dots$) are included self-consistently. The RMF functional can be one of the following four forms: the meson exchange or point-coupling nucleon interactions combined with the non-linear or density-dependent couplings. We solve the Dirac equation in an axially deformed harmonic oscillator (ADHO) basis. The convergence of the calculated results against the basis truncation and the basis deformation is studied and it is shown that reasonably large ADHO basis is able to provide a desired accuracy.

Three-dimensional potential energy surface in the (β_{20} , β_{22} , β_{30}) plane is obtained for ^{240}Pu and potential energy curves of actinide nuclei around the first and second fission barriers are studied systematically. It is found that the triaxiality is crucial in determining the height of both the first and the second fission barriers. However, most of actinides studied here are axially deformed at the first and second saddle points. The triaxial shape is mostly important before the saddle point around the first barrier, while it is often so after the saddle point around the

second barrier. Taking ^{234}U as an example, we have studied the parameter dependence of the effects of triaxiality on the second barrier height and found that the height of the second barrier may differ by a few MeV with different effective interactions, but in all cases the barriers are lowered considerably by the non-axial deformations.

We conclude that it is important to include the reflection asymmetric and non-axial shapes simultaneously for the study of potential energy surfaces and fission barriers of actinide nuclei and of those in unknown mass regions such as superheavy nuclei.

ACKNOWLEDGMENTS

Helpful discussions with Jie Meng, P. Ring, D. Vretnar, Xi-Zhen Wu, and Zhen-Hua Zhang are acknowledged. This work has been supported by Major State Basic Research Development Program of China (Grant No. 2013CB834400), National Natural Science Foundation of China (Grant Nos. 11121403, 11175252, 11120101005, 11211120152, and 11275248), Knowledge Innovation Project of Chinese Academy of Sciences (Grant Nos. KJCX2-EW-N01 and KJCX2-YW-N32). The results described in this paper are obtained on the ScGrid of Supercomputing Center, Computer Network Information Center of Chinese Academy of Sciences.

Appendix A: Matrix elements of the Dirac Hamiltonian

In the ADHO basis, the Dirac equation (5) is transformed into a matrix eigenvalue problem,

$$\begin{aligned} \sum_{\alpha'} \langle \alpha | M_+ (\mathbf{r}) | \alpha' \rangle f_i^{\alpha'} + \sum_{\alpha'} \langle \alpha | \boldsymbol{\sigma} \cdot \mathbf{p} | \alpha' \rangle g_i^{\alpha'} &= \epsilon_i f_i^\alpha, \\ \sum_{\alpha'} \langle \alpha | \boldsymbol{\sigma} \cdot \mathbf{p} | \alpha' \rangle f_i^{\alpha'} + \sum_{\alpha'} \langle \alpha | M_- (\mathbf{r}) | \alpha' \rangle g_i^{\alpha'} &= \epsilon_i g_i^\alpha. \end{aligned}$$

In the cylindrical coordinate system the kinetic energy operator $\boldsymbol{\sigma} \cdot \mathbf{p}$ reads,

$$\boldsymbol{\sigma} \cdot \mathbf{p} = -i \begin{pmatrix} \partial_z & e^{-i\varphi}(\partial_\rho - i\rho^{-1}\partial_\varphi) \\ e^{i\varphi}(\partial_\rho + i\rho^{-1}\partial_\varphi) & -\partial_z \end{pmatrix}.$$

The matrix elements of the kinetic energy operator $\boldsymbol{\sigma} \cdot \mathbf{p}$ can be calculated as:

$$\begin{aligned} \langle \alpha | \boldsymbol{\sigma} \cdot \mathbf{p} | \alpha' \rangle &= \langle n_z, n_\rho, m_l, m_s | \boldsymbol{\sigma} \cdot \mathbf{p} | n'_z, n'_\rho, m'_l, m'_s \rangle \\ &= -i C_\alpha^* C_{\alpha'} \delta_{K, K'} \left[\delta_{m_l, m'_l} \delta_{n_\rho, n'_\rho} \delta_{m_s, m'_s} A_{\alpha\alpha'} \right. \\ &\quad + \delta_{m_s, -\frac{1}{2}} \delta_{m'_s, \frac{1}{2}} \delta_{n_z, n'_z} (B_{\alpha\alpha'} - m'_l C_{\alpha\alpha'}) \\ &\quad \left. + \delta_{m_s, \frac{1}{2}} \delta_{m'_s, -\frac{1}{2}} \delta_{n_z, n'_z} (B_{\alpha\alpha'} + m'_l C_{\alpha\alpha'}) \right], \end{aligned} \quad (\text{A1})$$

where the integrals read,

$$A_{\alpha\alpha'} = \int dz \phi_{n_z} \partial_z \phi_{n'_z}, \quad (\text{A2})$$

$$B_{\alpha\alpha'} = \int d\rho R_{n_\rho}^{m_l} \rho \partial_\rho R_{n_\rho}^{m'_l}, \quad (\text{A3})$$

$$C_{\alpha\alpha'} = \int d\rho R_{n_\rho}^{m_l} R_{n_\rho}^{m'_l}. \quad (\text{A4})$$

The matrix elements of the potentials $U(\mathbf{r}) = M_\pm(\mathbf{r})$ are calculated by expanding $U(\mathbf{r})$ into a Fourier series,

$$U(\mathbf{r}) = U(z, \rho, \varphi) = \frac{1}{2\pi} \sum_{\mu=-\infty}^{\infty} U^{(\mu)}(z, \rho) e^{i\mu\varphi}, \quad (\text{A5})$$

where the components $U^{(\mu)}(z, \rho)$ are calculated by the inverse Fourier transformation,

$$U^{(\mu)}(z, \rho) = \int_0^{2\pi} d\varphi U(z, \rho, \varphi) e^{-i\mu\varphi}, \quad (\text{A6})$$

where the integral over φ is performed on a uniformly distributed mesh points. From the symmetry conditions it is easy to find,

$$U^{(\mu)} = U^{(-\mu)} = (-1)^\mu U^{(-\mu)}. \quad (\text{A7})$$

Thus the Fourier series is abbreviated as

$$U(\mathbf{r}) = \frac{1}{2\pi} \left(U^{(0)}(z, \rho) + 2 \sum_{n=1}^{\infty} U^{(2n)}(z, \rho) \cos(2n\varphi) \right). \quad (\text{A8})$$

Finally the matrix element of $U(\mathbf{r})$ reads,

$$\begin{aligned} \langle \alpha | U | \alpha' \rangle &= \langle n_z, n_r, m_l, m_s | U | n'_z, n'_r, m'_l, m'_s \rangle \\ &= \delta_{m_s, m'_s} C_\alpha^* C_{\alpha'} \frac{1}{2\pi} \left[\delta_{K, K'} D_{\alpha\alpha'}^{(0)} \right. \\ &\quad \left. + \sum_{n=1}^{\infty} (\delta_{K'-K, 2n} + \delta_{K-K', 2n}) D_{\alpha\alpha'}^{(2n)} \right] \end{aligned} \quad (\text{A9})$$

where we have performed the integrals over φ and

$$D_{\alpha\alpha'}^{(\mu)} = \int_{-\infty}^{\infty} dz \int_0^{\infty} \rho d\rho U^{(\mu)}(z, \rho) \phi_{n_z} \phi_{n'_z} R_{n_\rho}^{m_l} R_{n'_\rho}^{m'_l}, \quad (\text{A10})$$

are performed numerically by using the Gaussian quadrature. For axial symmetric potentials only the first term survives, while for triaxially deformed nucleus we must calculate additional terms with $n \neq 0$.

Appendix B: Densities and their derivatives

After solving the Dirac equation, we can calculate the densities from f_i^α and g_i^α . All the densities are linear combinations of the quantities $\rho_f(\mathbf{r}, \tau)$ and $\rho_g(\mathbf{r}, \tau)$ which are the contributions to the vector density from the large and small components, respectively. τ represents the isospin. For vector potential we have

$$\rho_V(\mathbf{r}) = \sum_{i=1}^N v_i^2 \sum_{p=f,g} \psi_i^{p\dagger} \psi_i^p$$

$$= \sum_{p=f,g} \sum_{\alpha\alpha'} \left(\sum_{i=1}^N v_i^2 p_i^\alpha p_i^{\alpha'} \right) \Phi_\alpha^\dagger \Phi_{\alpha'}. \quad (\text{B1})$$

Note that the factors with φ in Φ_α and $\Phi_{\alpha'}$ are in an exponential form, rearranging the terms, $\rho_V(\mathbf{r})$ can be written as,

$$\rho_V(\mathbf{r}) = \frac{1}{2\pi} \left(\rho_V^{(0)}(z, \rho) + 2 \sum_{n=1}^{\infty} \rho_V^{(2n)}(z, \rho) \cos(2n\varphi) \right), \quad (\text{B2})$$

where the components

$$\begin{aligned} \rho_V^{(\mu)} &= \sum_{p=f,g} \sum_{\alpha\alpha'} \left[\left(\sum_{i=1}^N v_i^2 p_i^\alpha p_i^{\alpha'} \right) \delta_{K'-K, \mu} \delta_{m_s, m'_s} \right. \\ &\quad \left. \times C_\alpha^* C_{\alpha'} \phi_{n_z} \phi_{n'_z} R_{n_\rho}^{m_l} R_{n'_\rho}^{m'_l} \right], \end{aligned} \quad (\text{B3})$$

are calculated with the wave functions f_i^α and g_i^α .

The scalar density $\rho_S(\mathbf{r})$ can be calculated similarly. The isovector densities are just the linear combinations of the corresponding densities of protons and neutrons.

The derivatives of the vector density reads

$$\nabla^2 \rho_V(\mathbf{r}) = \sum_{p=f,g} \sum_{\alpha\alpha'} \left(\sum_{i=1}^N v_i^2 p_i^\alpha p_i^{\alpha'} \right) \nabla^2 (\Phi_\alpha^\dagger \Phi_{\alpha'}), \quad (\text{B4})$$

where the derivatives of the basis can be calculated as,

$$\nabla^2 (\Phi_\alpha^\dagger \Phi_{\alpha'}) = I_{\alpha\alpha'} + 2J_{\alpha\alpha'} + I_{\alpha'\alpha}^*, \quad (\text{B5})$$

with $I_{\alpha\alpha'}$ and $J_{\alpha\alpha'}$ defined later. First, from the eigen equation Eq. (10) we get

$$\nabla^2 \Phi_\alpha = -2M [E_\alpha - V_B(z, \rho)] \Phi_\alpha, \quad (\text{B6})$$

thus

$$\begin{aligned} I_{\alpha\alpha'} &\equiv \Phi_\alpha^\dagger \nabla^2 \Phi_{\alpha'} \\ &= -2M [E_\alpha - V_B(z, \rho)] \delta_{m_s, m'_s} C_\alpha^* C_{\alpha'} \\ &\quad \times \phi_{n_z} \phi_{n'_z} R_{n_\rho}^{m_l} R_{n'_\rho}^{m'_l} \times \frac{1}{2\pi} e^{i(m'_l - m_l)\varphi}. \end{aligned} \quad (\text{B7})$$

Second,

$$\begin{aligned} J_{\alpha\alpha'} &\equiv \nabla \Phi_\alpha^\dagger \cdot \nabla \Phi_{\alpha'} \\ &= \delta_{m_s, m'_s} C_\alpha^* C_{\alpha'} \\ &\quad \times \left[\phi_{n_z} \phi_{n'_z} \left(\partial_\rho R_{n_\rho}^{m_l} \partial_\rho R_{n'_\rho}^{m'_l} + m_l m'_l \rho^{-2} R_{n_\rho}^{m_l} R_{n'_\rho}^{m'_l} \right) \right. \\ &\quad \left. + R_{n_\rho}^{m_l} R_{n'_\rho}^{m'_l} \partial_z \phi_{n_z} \partial_z \phi_{n'_z} \right] \times \frac{1}{2\pi} e^{i(m'_l - m_l)\varphi}. \end{aligned} \quad (\text{B8})$$

Substituting them into Eq. (B5) and combining the same exponentials of φ , we get

$$\nabla^2 \rho_V(\mathbf{r}) = \frac{1}{2\pi} \left(\tilde{\rho}_V^{(0)}(z, \rho) + 2 \sum_{n=1}^{\infty} \tilde{\rho}_V^{(2n)}(z, \rho) \cos(2n\varphi) \right), \quad (\text{B9})$$

in which

$$\begin{aligned} \tilde{\rho}_V^{(\mu)} = & \sum_{p=f,g} \sum_{\alpha\alpha'} \left(\sum_{i=1}^N v_i^2 p_i^\alpha p_i^{\alpha'} \right) \delta_{K'-K,\mu} \delta_{m_s,m'_s} \\ & \times C_{\alpha}^* C_{\alpha'} \left\{ D_{\alpha\alpha'}(z, \rho) \phi_{n_z} \phi_{n'_z} R_{n_\rho}^{m_l} R_{n'_\rho}^{m'_l} \right. \end{aligned}$$

$$\left. + 2\phi_{n_z} \phi_{n'_z} \partial_\rho R_{n_\rho}^{m_l} \partial_\rho R_{n'_\rho}^{m'_l} + 2R_{n_\rho}^{m_l} R_{n'_\rho}^{m'_l} \partial_z \phi_{n_z} \partial_z \phi_{n'_z} \right\}, \quad (\text{B10})$$

are the Fourier components and

$$D_{\alpha\alpha'}(z, \rho) = \frac{2m_l m'_l}{\rho^2} - 2M [E_{\alpha'} + E_\alpha - 2V_B(z, \rho)]. \quad (\text{B11})$$

-
- [1] A. Bohr and B. R. Mottelson, *Nuclear Structure*, Vol. I (World Scientific, 1998).
- [2] P. Ring and P. Schuck, *The Nuclear Many-Body Problem* (Springer-Verlag Berlin, Heidelberg, and New York, 1980).
- [3] S. Frauendorf, *Rev. Mod. Phys.* **73**, 463 (2001).
- [4] W. Nazarewicz, in *Lecture Notes in Physics*, Vol. 581, edited by J. Arias and M. Lozano (Springer Berlin Heidelberg, 2001) pp. 102–140.
- [5] F. Iachello, *Phys. Rev. Lett.* **87**, 052502 (2001).
- [6] Z. P. Li, T. Niksic, D. Vretenar, J. Meng, G. A. Lalazissis, and P. Ring, *Phys. Rev. C* **79**, 054301 (2009).
- [7] Z. P. Li, T. Niksic, D. Vretenar, and J. Meng, *Phys. Rev. C* **80**, 061301(R) (2009).
- [8] P. Cejnar, J. Jolie, and R. F. Casten, *Rev. Mod. Phys.* **82**, 2155 (2010).
- [9] J. Meng, W. Zhang, S. G. Zhou, H. Toki, and L. S. Geng, *Eur. Phys. J. A* **25**, 23 (2005), [arXiv:nucl-th/0312055](#).
- [10] R. Fossion, D. Bonatsos, and G. A. Lalazissis, *Phys. Rev. C* **73**, 044310 (2006).
- [11] J.-Y. Guo, X.-Z. Fang, and Z.-Q. Sheng, *Int. J. Mod. Phys. E* **17**, 539 (2008).
- [12] T. Niksic, D. Vretenar, G. A. Lalazissis, and P. Ring, *Phys. Rev. Lett.* **99**, 092502 (2007).
- [13] K. Nomura, N. Shimizu, D. Vretenar, T. Niksic, and T. Otsuka, *Phys. Rev. Lett.* **108**, 132501 (2012).
- [14] M. Brack, J. Damgaard, A. S. Jensen, H. C. Pauli, V. M. Strutinsky, and C. Y. Wong, *Rev. Mod. Phys.* **44**, 320 (1972).
- [15] R. W. Hasse and W. D. Myers, *Geometrical relationships of macroscopic nuclear physics*, Springer Series in Nuclear and Particle Physics (Springer-Verlag, 1988).
- [16] M. G. Mustafa, U. Mosel, and H. W. Schmitt, *Phys. Rev. Lett.* **28**, 1536 (1972).
- [17] W. J. Swiatecki, *Phys. Scr.* **24**, 113 (1981).
- [18] P. Möller, D. G. Madland, A. J. Sierk, and A. Iwamoto, *Nature* **409**, 785 (2001).
- [19] A. Diaz-Torres, *Phys. Rev. Lett.* **101**, 122501 (2008).
- [20] Q. Sun, D.-H. Shangguan, and J.-D. Bao, *Chin. Phys. C* **37**, 014102 (2013).
- [21] S. G. Nilsson, *Dan. Mat. Fys. Medd.* **29**, 16 (1955).
- [22] S. Cwiok, P.-H. Heenen, and W. Nazarewicz, *Nature* **433**, 705 (2005).
- [23] S. Frauendorf and J. Meng, *Nucl. Phys. A* **617**, 131 (1997).
- [24] K. Starosta, T. Koike, C. J. Chiara, D. B. Fossan, D. R. LaFosse, A. A. Hecht, C. W. Beausang, M. A. Caprio, J. R. Cooper, R. Krucken, J. R. Novak, N. V. Zamfir, K. E. Zyromski, D. J. Hartley, D. L. Balabanski, J.-y. Zhang, S. Frauendorf, and V. I. Dimitrov, *Phys. Rev. Lett.* **86**, 971 (2001).
- [25] S. W. Odegard, G. B. Hagemann, D. R. Jensen, M. Bergstroem, B. Herskind, G. Sletten, S. Toermannen, J. N. Wilson, P. O. Tjom, I. Hamamoto, K. Spohr, H. Huebel, A. Goergen, G. Schoenwasser, A. Bracco, S. Leoni, A. Maj, C. M. Petrache, P. Bednarczyk, and D. Curien, *Phys. Rev. Lett.* **86**, 5866 (2001).
- [26] J. Meng and S. Q. Zhang, *J. Phys. G: Nucl. Phys.* **37**, 064025 (2010).
- [27] J. Meng, J. Peng, S.-Q. Zhang, and P.-W. Zhao, *Frontiers of Physics* **8**, 55 (2013).
- [28] J. Meng, J. Peng, S. Q. Zhang, and S.-G. Zhou, *Phys. Rev. C* **73**, 037303 (2006), [arXiv:nucl-th/0510071](#).
- [29] A. D. Ayangeakaa, U. Garg, M. D. Anthony, S. Frauendorf, J. T. Matta, B. K. Nayak, D. Patel, Q. B. Chen, S. Q. Zhang, P. W. Zhao, B. Qi, J. Meng, R. V. F. Janssens, M. P. Carpenter, C. J. Chiara, F. G. Kondev, T. Lauritsen, D. Seweryniak, S. Zhu, S. S. Ghugre, and R. Palit, “Observation of multiple chiral doublet bands in ^{133}Ce ,” (2013), *Phys. Rev. Lett.*, in press, [arXiv:1302.0401 \[nucl-ex\]](#).
- [30] Z. P. Li, T. Niksic, D. Vretenar, and J. Meng, *Phys. Rev. C* **81**, 034316 (2010).
- [31] P. W. Zhao, J. Peng, H. Z. Liang, P. Ring, and J. Meng, *Phys. Rev. Lett.* **107**, 122501 (2011).
- [32] C.-Y. Song, Z.-P. Li, D. Vretenar, and J. Meng, *Sci. China-Phys. Mech. Astron.* **54**, 222 (2011).
- [33] Y. Zhang, Y.-X. Liu, Z.-F. Hou, and Y. Sun, *Sci. China-Phys. Mech. Astron.* **54**, 227 (2011).
- [34] S. Zhu, J. Wang, L. Gu, J. Hamilton, A. Ramayya, Y. Luo, J. Rasmussen, J. Hwang, H. Ding, K. Li, S. Liu, E. Yeoh, Q. Xu, and Z. Xiao, *Sci. China-Phys. Mech. Astron.* **54**, 44 (2011).
- [35] L. Liu and P. Zhao, *Sci. China-Phys. Mech. Astron.* **55**, 2420 (2012).
- [36] P. A. Butler and W. Nazarewicz, *Rev. Mod. Phys.* **68**, 349 (1996).
- [37] T. M. Shneidman, G. G. Adamian, N. V. Antonenko, R. V. Jolos, and W. Scheid, *Phys. Rev. C* **67**, 014313 (2003).
- [38] T. M. Shneidman, G. G. Adamian, N. V. Antonenko, and R. V. Jolos, *Phys. Rev. C* **74**, 034316 (2006).
- [39] S. Wang, H. Hua, J. Meng, Z. H. Li, S. Q. Zhang, F. R. Xu, H. L. Liu, Y. L. Ye, D. X. Jiang, T. Zheng, Q. J. Wang, Z. Q. Chen, C. E. Wu, G. L. Zhang, D. Y. Pang, J. Wang, J. L. Lou, B. Guo, G. Jin, S. G. Zhou, L. H. Zhu, X. G. Wu, G. S. Li, S. X. Wen, C. Y. He, X. Z. Cui, and Y. Liu, *Phys. Rev. C* **72**, 024317 (2005).
- [40] D. Yang, J.-B. Lu, Y.-Z. Liu, L.-L. Wang, K.-Y. Ma, C.-D. Yang, D.-K. Han, Y.-X. Zhao, Y.-J. Ma, L.-H. Zhu, X.-G. Wu, and G.-S. Li, *Chin. Phys. Lett.* **26**, 082101

- (2009).
- [41] L. M. Robledo and G. F. Bertsch, *Phys. Rev. C* **84**, 054302 (2011).
 - [42] S. J. Zhu, M. Sakhaee, J. H. Hamilton, A. V. Ramayya, N. T. Brewer, J. K. Hwang, S. H. Liu, E. Y. Yeoh, Z. G. Xiao, Q. Xu, Z. Zhang, Y. X. Luo, J. O. Rasmussen, I. Y. Lee, K. Li, and W. C. Ma, *Phys. Rev. C* **85**, 014330 (2012).
 - [43] S. J. Zhu, J. H. Hamilton, A. V. Ramayya, J. K. Hwang, Y. J. Chen, L. Y. Zhu, H. J. Li, Z. G. Xiao, E. Y. Yeoh, J. G. Wang, Y. X. Luo, S. H. Liu, J. O. Rasmussen, and I. Y. Lee, in *Nuclear Structure in China 2012: Proceedings of the 14th National Conference on Nuclear Structure in China (NSC2012)*, edited by J. Meng, C.-W. Shen, E.-G. Zhao, and S.-G. Zhou (2012) pp. 348–356.
 - [44] L. F. Yu, P. W. Zhao, S. Q. Zhang, and J. Meng, “Octupole deformation and Ra puzzle in reflection asymmetric covariant density functional theory,” arXiv:1211.0601 [nucl-th] (2012).
 - [45] D. S. Brenner, R. B. Cakirli, and R. F. Casten, *Phys. Rev. C* **73**, 034315 (2006).
 - [46] D. Neidherr, G. Audi, D. Beck, K. Blaum, C. Böhm, M. Breitenfeldt, R. B. Cakirli, R. F. Casten, S. George, F. Herfurth, A. Herlert, A. Kellerbauer, M. Kowalska, D. Lunney, E. Minaya-Ramirez, S. Naimi, E. Noah, L. Penescu, M. Rosenbusch, S. Schwarz, L. Schweikhard, and T. Stora, *Phys. Rev. Lett.* **102**, 112501 (2009).
 - [47] I. Hamamoto, B. Mottelson, H. Xie, and X. Z. Zhang, *Z. Phys. D* **21**, 163 (1991).
 - [48] J. Skalski, *Phys. Rev. C* **43**, 140 (1991).
 - [49] X. Li and J. Dudek, *Phys. Rev. C* **49**, R1250 (1994).
 - [50] S. Takami, K. Yabana, and M. Matsuo, *Phys. Lett. B* **431**, 242 (1998).
 - [51] M. Yamagami, K. Matsuyanagi, and M. Matsuo, *Nucl. Phys. A* **693**, 579 (2001).
 - [52] J. Dudek, A. Gozdz, N. Schunck, and M. Miskiewicz, *Phys. Rev. Lett.* **88**, 252502 (2002).
 - [53] J. Dudek, D. Curien, N. Dubray, J. Dobaczewski, V. Pangon, P. Olbratowski, and N. Schunck, *Phys. Rev. Lett.* **97**, 072501 (2006).
 - [54] P. Olbratowski, J. Dobaczewski, P. Powalowski, M. Sadziak, and K. Zborecki, *Int. J. Mod. Phys. E* **15**, 333 (2006).
 - [55] K. Zborecki, P. Magierski, P.-H. Heenen, and N. Schunck, *Phys. Rev. C* **74**, 051302(R) (2006).
 - [56] J. Dudek, A. Gozdz, K. Mazurek, and H. Molique, *J. Phys. G: Nucl. Phys.* **37**, 064032 (2010).
 - [57] A. P. Robinson, T. L. Khoo, I. Ahmad, S. K. Tandel, F. G. Kondev, T. Nakatsukasa, D. Seweryniak, M. Asai, B. B. Back, M. P. Carpenter, P. Chowdhury, C. N. Davids, S. Eeckhaudt, J. P. Greene, P. T. Greenlees, S. Gros, A. Heinz, R.-D. Herzberg, R. V. F. Janssens, G. D. Jones, T. Lauritsen, C. J. Lister, D. Peterson, J. Qian, U. S. Tandel, X. Wang, and S. Zhu, *Phys. Rev. C* **78**, 034308 (2008).
 - [58] Y.-S. Chen, Y. Sun, and Z.-C. Gao, *Phys. Rev. C* **77**, 061305(R) (2008).
 - [59] J. Zhao, B.-N. Lu, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **86**, 057304 (2012), arXiv:1209.6567 [nucl-th].
 - [60] R. A. Bark, J. F. Sharpey-Schafer, S. M. Maliage, T. E. Madiba, F. S. Komati, E. A. Lawrie, J. J. Lawrie, R. Lindsay, P. Maine, S. M. Mullins, S. H. T. Murray, N. J. Ncapayi, T. M. Ramashidza, F. D. Smit, and P. Vymers, *Phys. Rev. Lett.* **104**, 022501 (2010).
 - [61] M. Jentschel, W. Urban, J. Krempel, D. Tonev, J. Dudek, D. Curien, B. Lauss, G. de Angelis, and P. Petkov, *Phys. Rev. Lett.* **104**, 222502 (2010).
 - [62] S. G. Nilsson, C. F. Tsang, A. Sobczewski, Z. Szymaniński, S. Wycech, C. Gustafson, I.-L. Lamm, P. Möller, and B. Nilsson, *Nucl. Phys. A* **131**, 1 (1969).
 - [63] H. L. Liu, F. R. Xu, and P. M. Walker, *Phys. Rev. C* **86**, 011301(R) (2012).
 - [64] Z.-H. Zhang, J. Meng, E.-G. Zhao, and S.-G. Zhou, “Rotational properties of the superheavy nucleus ^{256}Rf in particle-number conserving cranked shell model,” arXiv:1208.1156 [nucl-th] (2012).
 - [65] O. Hahn and F. Strassmann, *Naturwissenschaften*, *Naturwissenschaften* **27**, 11 (1939).
 - [66] N. Bohr and J. A. Wheeler, *Phys. Rev.* **56**, 426 (1939).
 - [67] A. Zubov, G. Adamian, and N. Antonenko, *Phys. Part. Nucl.* **40**, 847 (2009).
 - [68] C.-J. Xia, B.-X. Sun, E.-G. Zhao, and S.-G. Zhou, *Sci. China-Phys. Mech. Astron.* **54** (Suppl. 1), s109 (2011), arXiv:1101.2725 [nucl-th].
 - [69] S. Hofmann and G. Münzenberg, *Rev. Mod. Phys.* **72**, 733 (2000).
 - [70] Y. Oganessian, *J. Phys. G: Nucl. Phys.* **34**, R165 (2007); Y. T. Oganessian, F. S. Abdullin, P. D. Bailey, D. E. Benker, M. E. Bennett, S. N. Dmitriev, J. G. Ezold, J. H. Hamilton, R. A. Henderson, M. G. Itkis, Y. V. Lobanov, A. N. Mezentssev, K. J. Moody, S. L. Nelson, A. N. Polyakov, C. E. Porter, A. V. Ramayya, F. D. Riley, J. B. Roberto, M. A. Ryabinin, K. P. Rykaczewski, R. N. Sagaidak, D. A. Shaughnessy, I. V. Shirokovsky, M. A. Stoyer, V. G. Subbotin, R. Sudowe, A. M. Sukhov, Y. S. Tsyganov, V. K. Utyonkov, A. A. Voinov, G. K. Vostokin, and P. A. Wilk, *Phys. Rev. Lett.* **104**, 142502 (2010).
 - [71] K. Morita, K. Morimoto, D. Kaji, T. Akiyama, S.-i. Goto, H. Haba, E. Ideguchi, R. Kanungo, K. Katori, H. Koura, H. Kudo, T. Ohnishi, A. Ozawa, T. Suda, K. Sueki, H.-S. Xu, T. Yamaguchi, A. Yoneda, A. Yoshida, and Y.-L. Zhao, *J. Phys. Soc. Jpn.* **73**, 2593 (2004); K. Morita, K. Morimoto, D. Kaji, H. Haba, K. Ozeki, Y. Kudou, T. Sumita, Y. Wakabayashi, A. Yoneda, K. Tanaka, S. Yamaki, R. Sakai, T. Akiyama, S.-i. Goto, H. Hasebe, M. Huang, T. Huang, E. Ideguchi, Y. Kasamatsu, K. Katori, Y. Kariya, H. Kikunaga, H. Koura, H. Kudo, A. Mashiko, K. Mayama, S.-i. Mitsuoka, T. Moriya, M. Murakami, H. Murayama, S. Namai, A. Ozawa, N. Sato, K. Sueki, M. Takeyama, F. Tokanai, T. Yamaguchi, and A. Yoshida, *J. Phys. Soc. Jpn.* **81**, 103201 (2012).
 - [72] Z.-Y. Zhang, Z.-G. Gan, L. Ma, M.-H. Huang, T.-H. Huang, X.-L. Wu, G.-B. Jia, G.-S. Li, Z.-Z. Ren, S.-G. Zhou, X.-H. Zhou, H.-S. Xu, H.-Q. Zhang, and G.-Q. Xiao, *Chin. Phys. Lett.* **29**, 012502 (2012).
 - [73] P. Möller, A. J. Sierk, T. Ichikawa, A. Iwamoto, R. Bengtsson, H. Uhlenholt, and S. Aberg, *Phys. Rev. C* **79**, 064304 (2009).
 - [74] J. Peter, *Eur. Phys. J. A* **22**, 271 (2004); *Nucl. Phys. A* **734**, 192 (2004).
 - [75] J. C. Pei, W. Nazarewicz, J. A. Sheikh, and A. K. Kerman, *Phys. Rev. Lett.* **102**, 192501 (2009).
 - [76] N. V. Antonenko, E. A. Cherepanov, A. K. Nasirov, V. P. Permjakov, and V. V. Volkov, *Phys. Rev. C* **51**,

- 2635 (1995).
- [77] G. G. Adamian, N. V. Antonenko, and W. Scheid, *Nucl. Phys. A* **618**, 176 (1997).
 - [78] Z. H. Liu and J.-D. Bao, *Phys. Rev. C* **76**, 034604 (2007); Z.-H. Liu and J.-D. Bao, *Phys. Rev. C* **87**, 034616 (2013).
 - [79] Z.-G. Gan, X.-H. Zhou, M.-H. Huang, Z.-Q. Feng, and J.-Q. Li, *Sci. China-Phys. Mech. Astron.* **54** (Supp. 1), s61 (2011).
 - [80] N. Wang, J. Tian, and W. Scheid, *Phys. Rev. C* **84**, 061601(R) (2011); V. Zanganeh, N. Wang, and O. N. Ghodsi, *Phys. Rev. C* **85**, 034601 (2012).
 - [81] P. M. Walker, F. R. Xu, H. L. Liu, and Y. Sun, *J. Phys. G: Nucl. Part. Phys.* **39**, 105106 (2012).
 - [82] P. Möller, A. J. Sierk, and A. Iwamoto, *Phys. Rev. Lett.* **92**, 072501 (2004).
 - [83] K. Pomorski and J. Dudek, *Phys. Rev. C* **67**, 044316 (2003).
 - [84] F. A. Ivanyuk and K. Pomorski, *Phys. Rev. C* **79**, 054327 (2009).
 - [85] M. Kowal, P. Jachimowicz, and A. Sobiczewski, *Phys. Rev. C* **82**, 014303 (2010).
 - [86] G. Royer, M. Jaffre, and D. Moreau, *Phys. Rev. C* **86**, 044326 (2012).
 - [87] A. Mamdouh, J. M. Pearson, M. Rayet, and F. Tondeur, *Nucl. Phys. A* **644**, 389 (1998); *Nucl. Phys. A* **679**, 337 (2001).
 - [88] T. Burvenich, M. Bender, J. A. Maruhn, and P.-G. Reinhard, *Phys. Rev. C* **69**, 014307 (2004).
 - [89] M. Samyn, S. Goriely, and J. M. Pearson, *Phys. Rev. C* **72**, 044316 (2005); S. Goriely, M. Samyn, and J. M. Pearson, *Phys. Rev. C* **75**, 064312 (2007).
 - [90] F. Minato, S. Chiba, and K. Hagino, *Nucl. Phys. A* **831**, 150 (2009).
 - [91] M. Kortelainen, J. McDonnell, W. Nazarewicz, P.-G. Reinhard, J. Sarich, N. Schunck, M. V. Stoitsov, and S. M. Wild, *Phys. Rev. C* **85**, 024304 (2012).
 - [92] J. McDonnell, N. Schunck, and W. Nazarewicz, “Microscopic description of nuclear fission: Fission barrier heights of even-even actinides,” arXiv:1301.7587 [nucl-th] (2013), Proceedings of the 5th International Conference on Fission and properties of neutron-rich nuclei (ICFN5), Sanibel Island, Nov. 4-10, 2012.
 - [93] A. Staszczak, A. Baran, and W. Nazarewicz, *Phys. Rev. C* **87**, 024320 (2013).
 - [94] J. L. Egido and L. M. Robledo, *Phys. Rev. Lett.* **85**, 1198 (2000).
 - [95] M. Warda and J. L. Egido, *Phys. Rev. C* **86**, 014322 (2012).
 - [96] V. Blum, J. Maruhn, P.-G. Reinhard, and W. Greiner, *Phys. Lett. B* **323**, 262 (1994).
 - [97] W. Zhang, S.-S. Zhang, S.-Q. Zhang, and J. Meng, *Chin. Phys. Lett.* **20**, 1694 (2003).
 - [98] H.-F. Lü, L.-S. Geng, and J. Meng, *Chin. Phys. Lett.* **23**, 2940 (2006).
 - [99] Z. P. Li, T. Niksic, D. Vretenar, P. Ring, and J. Meng, *Phys. Rev. C* **81**, 064321 (2010).
 - [100] H. Abusara, A. V. Afanasjev, and P. Ring, *Phys. Rev. C* **82**, 044303 (2010).
 - [101] H. Abusara, A. V. Afanasjev, and P. Ring, *Phys. Rev. C* **85**, 024314 (2012).
 - [102] A. V. Afanasjev, “Fission of actinides and superheavy nuclei: covariant density functional theory perspective,” arXiv:1303.1206 [nucl-th] (2013), Proceedings of Fifth International conference on Fission and properties of neutron-rich nuclei, November 4-10, 2012, Sanibel Island, USA.
 - [103] R. Bengtsson, I. Ragnarsson, S. Aberg, A. Gyurkovich, A. Sobiczewski, and K. Pomorski, *Nucl. Phys. A* **473**, 77 (1987).
 - [104] P. Thirolf and D. Habs, *Prog. Part. Nucl. Phys.* **49**, 325 (2002).
 - [105] L. Csige, M. Csatlos, T. Faestermann, J. Gulyas, D. Habs, R. Hertenberger, M. Hunyadi, A. Krasznahorkay, H. J. Maier, P. G. Thirolf, and H.-F. Wirth, *Phys. Rev. C* **85**, 054306 (2012).
 - [106] M. Kowal and J. Skalski, *Phys. Rev. C* **85**, 061302(R) (2012).
 - [107] J. Jachimowicz, M. Kowal, and J. Skalski, “8Dim calculations of the third barrier in ^{232}Th and a conflict between theory and experiment on uranium nuclei,” arXiv:1301.0713 [nucl-th] (2013).
 - [108] V. V. Pashkevich, *Nucl. Phys. A* **133**, 400 (1969).
 - [109] P. Möller and S. G. Nilsson, *Phys. Lett. B* **31**, 283 (1970).
 - [110] J. Randrup, S. E. Larsson, P. Moller, S. G. Nilsson, K. Pomorski, and A. Sobiczewski, *Phys. Rev. C* **13**, 229 (1976).
 - [111] T. Ledergerber and H.-C. Pauli, *Nucl. Phys. A* **207**, 1 (1973).
 - [112] M. Girod and B. Grammaticos, *Phys. Rev. C* **27**, 2317 (1983).
 - [113] K. Rutz, J. A. Maruhn, P. G. Reinhard, and W. Greiner, *Nucl. Phys. A* **590**, 680 (1995).
 - [114] L. Bonneau, P. Quentin, and D. Samsen, *Eur. Phys. J. A* **21**, 391 (2004).
 - [115] P. Jachimowicz, M. Kowal, and J. Skalski, *Phys. Rev. C* **83**, 054302 (2011).
 - [116] J. Skalski, *Phys. Rev. C* **76**, 044603 (2007).
 - [117] B. D. Serot and J. D. Walecka, *Adv. Nucl. Phys.* **16**, 1 (1986).
 - [118] P. G. Reinhard, *Rep. Prog. Phys.* **52**, 439 (1989).
 - [119] P. Ring, *Prog. Part. Nucl. Phys.* **37**, 193 (1996).
 - [120] M. Bender, P.-H. Heenen, and P.-G. Reinhard, *Rev. Mod. Phys.* **75**, 121 (2003).
 - [121] D. Vretenar, A. Afanasjev, G. Lalazissis, and P. Ring, *Phys. Rep.* **409**, 101 (2005).
 - [122] J. Meng, H. Toki, S. G. Zhou, S. Q. Zhang, W. H. Long, and L. S. Geng, *Prog. Part. Nucl. Phys.* **57**, 470 (2006), arXiv:nucl-th/0508020.
 - [123] N. Paar, D. Vretenar, and G. Colo, *Rep. Prog. Phys.* **70**, 691 (2007).
 - [124] T. Niksic, D. Vretenar, and P. Ring, *Prog. Part. Nucl. Phys.* **66**, 519 (2011).
 - [125] B.-N. Lu, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **85**, 011301(R) (2012), arXiv:1110.6769 [nucl-th].
 - [126] B.-N. Lu, *Multi-dimensional constrained relativistic mean field theory and the potential energy surfaces and fission barriers of actinide nuclei*, Ph.D. thesis, Institute of Theoretical Physics, Chinese Academy of Sciences (2012), in Chinese.
 - [127] B.-N. Lu, J. Zhao, E.-G. Zhao, and S.-G. Zhou, *EPJ Web of Conferences* **38**, 05003 (2012), Proceedings of the Int. Conf. on Nuclear Structure and Related Topics (NSRT12), Dubna, Russia, 2-7 July 2012, arXiv:1303.0621 [nucl-th].
 - [128] V. Prassa, T. Niksic, G. A. Lalazissis, and D. Vretenar, *Phys. Rev. C* **86**, 024317 (2012).

- [129] Y. Tian and Z.-Y. Ma, *Chin. Phys. Lett.* **23**, 3226 (2006).
- [130] Y. Tian, Z. Ma, and P. Ring, *Phys. Lett. B* **676**, 44 (2009).
- [131] Y. Tian, Z.-y. Ma, and P. Ring, *Phys. Rev. C* **79**, 064301 (2009).
- [132] C. Fuchs, H. Lenske, and H. H. Wolter, *Phys. Rev. C* **52**, 3043 (1995).
- [133] T. Niksic, D. Vretenar, P. Finelli, and P. Ring, *Phys. Rev. C* **66**, 024306 (2002).
- [134] J. Boguta and A. R. Bodmer, *Nucl. Phys. A* **292**, 413 (1977).
- [135] R. Brockmann and H. Toki, *Phys. Rev. Lett.* **68**, 3408 (1992).
- [136] Y. Sugahara and H. Toki, *Nucl. Phys. A* **579**, 557 (1994).
- [137] Y. K. Gambhir, P. Ring, and A. Thimet, *Ann. Phys.* **198**, 132 (1990).
- [138] P. Ring, Y. K. Gambhir, and G. A. Lalazissis, *Comput. Phys. Commun.* **105**, 77 (1997).
- [139] S.-G. Zhou, J. Meng, and P. Ring, *Phys. Rev. C* **68**, 034323 (2003), [arXiv:nucl-th/0303031](#).
- [140] S.-G. Zhou, J. Meng, and P. Ring, *Nuclear Physics Trends, AIP Conf. Proc.*, **865**, 90 (2006).
- [141] S.-G. Zhou, J. Meng, P. Ring, and E.-G. Zhao, *Phys. Rev. C* **82**, 011301(R) (2010), [arXiv:0909.1600 \[nucl-th\]](#).
- [142] L. Li, J. Meng, P. Ring, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **85**, 024312 (2012), [arXiv:1202.0070 \[nucl-th\]](#).
- [143] Y. Chen, L. Li, H. Liang, and J. Meng, *Phys. Rev. C* **85**, 067301 (2012).
- [144] L.-S. Geng, J. Meng, and H. Toki, *Chin. Phys. Lett.* **24**, 1865 (2007).
- [145] B.-N. Lu, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **84**, 014328 (2011), [arXiv:1104.4638 \[nucl-th\]](#).
- [146] S. Karatzikos, A. Afanasjev, G. Lalazissis, and P. Ring, *Phys. Lett. B* **689**, 72 (2010).
- [147] J. Y. Zeng and T. S. Cheng, *Nucl. Phys. A* **405**, 1 (1983).
- [148] H. Molique and J. Dudek, *Phys. Rev. C* **56**, 1795 (1997).
- [149] J. Meng, J.-y. Guo, L. Liu, and S.-q. Zhang, *Frontiers Phys. China* **1**, 38 (2006).
- [150] N. Pillet, P. Quentin, and J. Libert, *Nucl. Phys. A* **697**, 141 (2002).
- [151] Z.-H. Zhang, J.-Y. Zeng, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **83**, 011304(R) (2011), [arXiv:1101.3607 \[nucl-th\]](#); Z.-H. Zhang, X.-T. He, J.-Y. Zeng, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **85**, 014324 (2012), [arXiv:1110.6134 \[nucl-th\]](#).
- [152] P. W. Zhao, Z. P. Li, J. M. Yao, and J. Meng, *Phys. Rev. C* **82**, 054319 (2010); P. W. Zhao, L. S. Song, B. Sun, H. Geissel, and J. Meng, *Phys. Rev. C* **86**, 064324 (2012).
- [153] I. Reichstein and F. Bary Malik, *Ann. Phys.* **98**, 322 (1976).
- [154] B. N. Pomorska, *Nucl. Phys. A* **327**, 1 (1979).
- [155] F. Tondeur, *Nucl. Phys. A* **442**, 460 (1985).
- [156] M. Bender, P.-H. Heenen, and P. Bonche, *Phys. Rev. C* **70**, 054304 (2004).
- [157] A. Andreev, G. Adamian, N. Antonenko, and S. Ivanova, *Eur. Phys. J. A* **26**, 327 (2005).
- [158] L. M. Robledo, M. Baldo, P. Schuck, and X. Vinas, *Phys. Rev. C* **77**, 051301 (2008).
- [159] W. Younes and D. Gogny, *Phys. Rev. C* **80**, 054313 (2009).
- [160] T. V. N. Hao, P. Quentin, and L. Bonneau, *Phys. Rev. C* **86**, 064307 (2012).
- [161] R. Capote, M. Herman, P. Oblozinsky, P. Young, S. Goriely, T. Belgia, A. Ignatyuk, A. Koning, S. Hilaire, V. Plujko, M. Avrigeanu, O. Bersillon, M. Chadwick, T. Fukahori, Z. Ge, Y. Han, S. Kailas, J. Kopecky, V. Maslov, G. Reffo, M. Sin, E. Soukhovitskii, and P. Talou, *Special Issue on Nuclear Reaction Data, Nucl. Data Sheets* **110**, 3107 (2009).
- [162] B.-N. Lu, J. Zhao, E.-G. Zhao, and S.-G. Zhou, (2013), Proceedings of the XXXVI Mexican Symposium on Nuclear Physics, Cocoyoc, Morelos, Mexico, Jan. 7-10, 2013.
- [163] G. A. Lalazissis, J. Konig, and P. Ring, *Phys. Rev. C* **55**, 540 (1997); G. A. Lalazissis, S. Karatzikos, R. Fossion, D. P. Arteaga, A. V. Afanasjev, and P. Ring, *Phys. Lett. B* **671**, 36 (2009).
- [164] M. Bender, K. Rutz, P.-G. Reinhard, J. A. Maruhn, and W. Greiner, *Phys. Rev. C* **60**, 034304 (1999).
- [165] G. A. Lalazissis, T. Niksic, D. Vretenar, and P. Ring, *Phys. Rev. C* **71**, 024312 (2005).
- [166] T. Niksic, D. Vretenar, and P. Ring, *Phys. Rev. C* **78**, 034318 (2008).
- [167] P. Jachimowicz, M. Kowal, and J. Skalski, *Phys. Rev. C* **85**, 034305 (2012).
- [168] N. Dubray and D. Regnier, *Comput. Phys. Commun.* **183**, 2035 (2012).
- [169] J. M. Yao, J. Meng, P. Ring, and D. Vretenar, *Phys. Rev. C* **81**, 044311 (2010); J. M. Yao, H. Mei, H. Chen, J. Meng, P. Ring, and D. Vretenar, *Phys. Rev. C* **83**, 014308 (2011); J.-M. Yao, J. Peng, J. Meng, and P. Ring, *Sci. China Phys. Mech. Astron.* **54**, 198 (2011).
- [170] H. L. Liu, F. R. Xu, Y. Sun, P. M. Walker, and R. Wyss, *Eur. Phys. J. A* **47**, 135 (2011).
- [171] T. V. Nhan Hao, J. Le Bloas, M.-H. Koh, L. Bonneau, and P. Quentin, *Int. J. Mod. Phys. E* **31**, 1250051 (2012).